

GUIDE TO INFECTION CONTROL IN THE HOSPITAL

CHAPTER 43:

Staphylococcus Aureus

Authors

J. Pierce, MD

M. Edmond, MD, MPH, MPA

M.P. Stevens, MD, MPH

Chapter Editor

Michelle Doll, MD, MPH)

Topic Outline

Key Issue

Known Facts

Controversial Issues

Suggested Practice

MSSA

MRSA/VISA

VRSA

Suggested Practice in Under-Resourced Settings

Summary

References

Chapter last updated: February, 2018

KEY ISSUES

Staphylococcus aureus is a major human pathogen that commonly causes healthcare-associated and community-acquired infections. It is a highly virulent organism that exhibits significant antibiotic resistance.

KNOWN FACTS

- Colonization with *S. aureus* is common. A national, population-based study of non-hospitalized persons in the U.S. found 32% of persons to be colonized with methicillin-susceptible *S. aureus* (MSSA) and 1% colonized with methicillin-resistant *S. aureus* (MRSA).
- *S. aureus* is a major cause of healthcare-associated infections (HAIs), accounting for 12% of all HAIs in the United States.
- Regarding antimicrobial resistance, *S. aureus* is typically characterized by its susceptibility patterns to penicillinase-resistant penicillins (e.g., methicillin) and vancomycin.
- The *mecA* gene encodes for penicillin binding protein 2a (PBP2a) which confers resistance to beta-lactam antibiotics.
- Over half of all *S. aureus* strains acquired in U.S. healthcare facilities are resistant to methicillin.
- Historically, MSSA strains were mostly acquired in the community, whereas MRSA strains were typically acquired in healthcare facilities.

However, community-associated MRSA (CA-MRSA) is now the predominant cause of purulent skin and soft tissue infections in the outpatient setting. MRSA isolates with the CA-MRSA phenotype are now commonly encountered in hospital settings, as well.

- Classification of MRSA strains into community-associated and hospital-associated based on exposure to the healthcare setting is no longer reliable.
- CA-MRSA tends to differ from traditional hospital-acquired MRSA in that community-associated strains are more likely to be susceptible to TMP/SMX and tetracyclines.
- CA-MRSA often manifests as skin and soft tissue infections and may be misdiagnosed as a “spider bite.” CA-MRSA is responsible for the majority of purulent skin and soft tissue infections presenting to U.S. emergency rooms.
- Many community-associated strains contain the Pantone-Valentine leukocidin (PVL) gene which is associated with lysis of white blood cells and tissue necrosis. These strains characteristically cause skin and soft tissue infections, often in healthy children and young adults, as well as a severe, multilobar, necrotizing pneumonia that often occurs with or following influenza.
- Risk factors for staphylococcal colonization and infection include disruptions of the skin (insulin injections, hemodialysis, allergy therapy, IV drug use, eczema, burns), underlying diseases (respiratory infections,

HIV infection), prolonged hospitalization, and exposure to other infected or colonized individuals. However, in many patients with CA-MRSA infections, these risk factors are not present.

- >80% of cases of *S. aureus* bacteremia are caused by endogenous strains (i.e., a strain colonizing the patient is responsible for invasive infection).
- The most common sources of *S. aureus* bloodstream infection are catheters (46%), skin/soft tissue/bone (27%), lower respiratory tract (11%), and urinary tract (10%).
- Vancomycin intermediate *S. aureus* (VISA), vancomycin resistant *S. aureus* (VRSA), and heteroresistant *S. aureus* (hetero-VRSA) have all been reported.
- The Clinical and Laboratory Standards Institute defines staphylococcal vancomycin minimum inhibitory concentrations (MICs) of ≤ 2 $\mu\text{g}/\text{mL}$ as susceptible, 4–8 $\mu\text{g}/\text{mL}$ as intermediate, and ≥ 16 $\mu\text{g}/\text{mL}$ as resistant. Generally speaking, vancomycin should be avoided for severe infections where the staphylococcal isolate has an MIC of ≥ 2 $\mu\text{g}/\text{mL}$ due to the risk of treatment failure.
- Hetero-VRSA are defined as strains of *S. aureus* that contain subpopulations of vancomycin-resistant daughter cells but for which the MICs of the parent strain are only 1–4 $\mu\text{g}/\text{mL}$. These subpopulations typically have MICs 2–8 fold higher than the original clinical isolate.

When grown in the absence of vancomycin, the subpopulation of cells reverts back to the lower MIC of the parent strain.

- In 2002, two strains of *S. aureus* with high levels of resistance to vancomycin (VRSA) were reported in the United States. These strains have MICs ≥ 16 $\mu\text{g/ml}$. As of February 2015, 14 patients in the U.S. had been identified with infections due to VRSA.
- Patients who develop infection with VISA and VRSA often have serious comorbid disease states such as renal failure and diabetes, a previous history of infections with MRSA, recent vancomycin use, the presence of foreign material (including intravenous catheters and prosthetic devices) and recent hospitalizations.
- Major route of transmission for *S. aureus* is direct or indirect contact; airborne transmission is uncommon.
- Colonized healthcare workers may be the source of outbreaks in the hospital setting.

Controversial Issues

- The effectiveness of routine surveillance cultures to detect MRSA colonized patients followed by isolation of the patient in order to reduce MRSA infection and colonization in high prevalence settings is probably not effective.

- The role of decolonizing agents in the non-outbreak setting remains undefined. In particular, use of mupirocin for all patients in the ICU setting (universal decolonization), raises concerns for the development of high rates of resistance. Resistance to chlorhexidine is also a concern (but appears to occur to a much less frequent extent).
- Use of contact precautions (gloves and gowns) in non-outbreak settings continues to be recommended by major organizations. However, increasing evidence (in resource rich environments) indicates that this may not be universally necessary.

SUGGESTED PRACTICE

MSSA

- Use standard precautions.

MRSA/VISA

- Use contact precautions (gloves and gowns).
- Emphasize handwashing with antiseptic agents (chlorhexidine gluconate or alcohol-based products).
- Consider private room or cohorting the infected or colonized patient with other MRSA patients.

- Offer decolonization with intranasal mupirocin and chlorhexidine bathing for patients with recurring infections and for colonized personnel.
- If the MRSA patient is transferred, notify the receiving healthcare facility.
- No special precautions for home discharge are required; emphasize good handwashing.
- Universal chlorhexidine bathing of ICU patients can have a major impact on reducing MRSA infections.

VRSA

- Contact precautions, including a private room, are recommended.
- Minimize the number of people in contact with or caring for the patient.
- Educate all healthcare personnel about the epidemiology of VRSA and the appropriate infection control precautions.
- Daily chlorhexidine bathing should be considered while an inpatient.
- Initiate epidemiologic and laboratory investigations with the assistance of the public health department.
- Consult with the public health department before transferring or discharging the patient.

SUGGESTED PRACTICE IN UNDER-RESOURCED SETTINGS

- In resource limited settings compliance with guidelines can be inconsistent; low nurse-to-patient staffing ratios, insufficient infection prevention training of health care workers, poor access to medical supplies, and hospital overcrowding can all contribute.
- Infection prevention training for healthcare workers is critical; strict compliance with handwashing should be emphasized.
- The other infection prevention strategies outlined above should be deployed whenever possible.

SUMMARY

- In the community, *S. aureus* is best known as the cause of furuncles and soft tissue infections. In the hospital environment, *S. aureus* may cause life-threatening infections, such as pneumonia, bloodstream, or surgical site infections, and is considered one of the most important hospital-acquired pathogens.
- The nares are the usual reservoir for *S. aureus*, but other locations such as moist or hairy body areas, skin defects, wounds, and burns also can become colonized. Methicillin-resistant *S. aureus* carriage may be

eradicated with application of topical mupirocin to the anterior nares, although recolonization often occurs. This therapy should be limited to patients with recurring MRSA infections, to select pre-operative patients who have documented nares MRSA colonization, and to colonized hospital personnel to prevent the development of resistance.

- The most common mode of *S. aureus* transmission is direct contact of body surface to body surface. Sexual transmission of MRSA has been described and manifests as folliculitis or abscesses of the pubic, vaginal or perineal areas. The airborne route is less efficient but may occur in patients with *S. aureus* pneumonia or large burn wounds. It has been shown that colonized individuals with viral upper respiratory tract infections may shed *S. aureus* into the air. Transmission via indirect contact with inanimate objects such as instruments can occur, and *S. aureus* can be detected on many surfaces in hospitals, including stethoscopes and laboratory coats.
- Strategies for the management of *S. aureus* and especially MRSA colonization or infection must focus on the type of spread. Epidemic outbreaks are successfully handled with prompt application of infection control measures. Application of precautions such as patient isolation, handwashing with antiseptic agents, and glove use can interrupt the chain of transmission and control the outbreak. Institutions with repeated introduction of MRSA from the community or other facilities are unlikely to be able to eradicate this pathogen.

- Vancomycin remains the mainstay of therapy for systemic MRSA infections. For MRSA-associated necrotizing pneumonia some experts recommend the addition of an antibiotic active at the ribosomal level (such as clindamycin) to terminate toxin production. For relatively minor skin infections, the use of doxycycline or trimethoprim/sulfamethoxazole (TMP/SMX) is typically recommended in addition to incision and drainage of abscesses.
- Fortunately, infections due to VISA and VRSA have remained uncommon. In the United States, there have been fourteen cases ascribed to VRSA. Importantly, strict compliance with infection control guidelines is necessary to minimize cross transmission within healthcare facilities. When identified, public health departments should be involved in the management of these cases.
- Treatment options for VISA and VRSA are few, and clinical experience is limited. Quinupristin-dalfopristin and linezolid are bacteriostatic for VISA/VRSA. Other potential therapies include daptomycin, ceftaroline, ceftobiprole, telavancin, tedizolid, and tigecycline. Susceptibility of VISA/VRSA has also been reported to chloramphenicol, minocycline, tetracycline, doxycycline, and TMP/SMX. Expert consultation with an infectious diseases specialist should be sought for the management of VISA and VRSA cases.

REFERENCES

1. Chang S, Sievert DM, Hageman JC, et al. Infection with Vancomycin-Resistant *Staphylococcus aureus* Containing the *vanA* Resistance Gene. *N Engl J Med* 2003; 348(14):1342–7.
2. Drew RH. Emerging Options for Treatment of Invasive, Multidrug-Resistant *Staphylococcus aureus* Infections. *Pharmacotherapy* 2007; 27(2):227–49.
3. Graham PL 3rd, Lin SX, Larson EL. A U.S. Population-Based Survey of *Staphylococcus aureus* Colonization. *Ann Intern Med* 2006; 144(5):318–25.
4. Liu C, Bayer A, Cosgrove SE, et al. Clinical Practice Guidelines by the Infectious Diseases Society of America for the Treatment of Methicillin-Resistant *Staphylococcus aureus* Infections in Adults and Children: Executive Summary. *Clin Infect Dis* 2011; 52(3):285–92. doi: 10.1093/cid/cir034.
5. Talan DA, Krishnadasan A, Gorwitz RJ, et al. Comparison of *Staphylococcus aureus* from Skin and Soft-Tissue Infections in US Emergency Department Patients, 2004 and 2008. *Clin Infect Dis* 2011; 53(2):144–9. doi: 10.1093/cid/cir308.

6. von Eiff C, Becker K, Machka K, et al. Nasal Carriage as a Source of *Staphylococcus aureus* Bacteremia Study Group. N Engl J Med 2001; 344(1):11–6.
7. Wenzel RP, Edmond MB. Vancomycin-Resistant *Staphylococcus aureus*: Infection Control Considerations. Clin Infect Dis 1998; 27(2):245–9, quiz 250-1.
8. Wisplinghoff H, Bischoff T, Tallent SM, et al. Nosocomial Bloodstream Infections in US Hospitals: Analysis of 24,179 Cases from a Prospective Nationwide Surveillance Study. Clin Infect Dis 2004; 39(3):309–17.
9. Weiner LM, Webb AK, Limbago B, et al. Antimicrobial-Resistant Pathogens Associated With Healthcare-Associated Infections: Summary of Data Reported to the National Healthcare Safety Network at the Centers for Disease Control and Prevention, 2011-2014. Infect Control Hosp Epidemiol 2016; 37(11):1288–301.
10. CDC. Antibiotic Resistance Threats in the United States, 2013; available at <http://www.cdc.gov/drugresistance/threat-report-2013>. Accessed 17 November 2017.

11. Huang SS, Septimus E, Kleinman K, et al. Targeted versus Universal Decolonization to Prevent ICU Infection. *N Engl J Med* 2013; 368(24):2255–65. doi: 10.1056/NEJMoa1207290.
12. Edmond MB, Masroor N, Stevens MP, et al. The Impact of Discontinuing Contact Precautions for VRE and MRSA on Device-Associated Infections. *Infect Control Hosp Epidemiol* 2015; 36(8):978–80. doi: 10.1017/ice.2015.99.
13. Rosenthal, VD, Al-Abdely, HM, El-Kholy, AA, et al. International Nosocomial Infection Control Consortium Report, Data Summary of 50 Countries for 2010-2015: Device-Associated Module. *Am J Infect Control*. 2016; 44(12):1495–1504. doi: 10.1016/j.ajic.2016.08.007.
14. Kampf G. Acquired Resistance to Chlorhexidine – Is It Time to Establish an “Antiseptic Stewardship” Initiative? *J Hosp Infect* 2016; 94(3):213-27. doi: 10.1016/j.jhin.2016.08.018